

**Chapter 7**

**Cell Structure and Function**

Lecture Presentations by  
Nicole Tunbridge and  
Kathleen Fitzpatrick

© 2018 Pearson Education Ltd.

## The Fundamental Units of Life


- All organisms are made of cells
- The cell is the simplest collection of matter that can be alive
- All cells are related by their descent from earlier cells
- Cells can differ substantially from one another but share common features

Figure 7.1



© 2018 Pearson Education Ltd.

Figure 7.1a



© 2018 Pearson Education Ltd.

**Concept 7.1: Biologists use microscopes and the tools of biochemistry to study cells**

- Cells are usually too small to be seen by the naked eye

© 2018 Pearson Education Ltd.

■ **Assignment:**


- Microscopes types & Functions
- Fractionation

- Three important parameters of microscopy:
  - Magnification, the ratio of an object's image size to its real size
  - Resolution, the measure of the clarity of the image, or the minimum distance of two distinguishable points
  - Contrast, visible differences in brightness between parts of the sample


© 2018 Pearson Education Ltd.

Figure 7.3b


### Light Microscopy (LM)




Fluorescence 10  $\mu\text{m}$



Deconvolution 10  $\mu\text{m}$




Confocal (without)



Confocal (with) 50  $\mu\text{m}$

© 2018 Pearson Education Ltd.



- Two basic types of **electron microscopes (EMs)** are used to study subcellular structures
- **Scanning electron microscopes (SEMs)** focus a beam of electrons onto the surface of a specimen, providing images that look 3-D
- **Transmission electron microscopes (TEMs)** focus a beam of electrons through a specimen
- TEMs are used mainly to study the internal structure of cells

- Recent advances in light microscopy:
  - Labeling individual cells with fluorescent markers improve the level of detail that can be seen
  - Confocal microscopy and deconvolution microscopy provide sharper images of three-dimensional tissues and cells
  - New techniques for labeling cells improve resolution
  - Super-resolution microscopy allows one to distinguish structures as small as 10–20 nm across

© 2018 Pearson Education Ltd.

### **Concept 7.2: Eukaryotic cells have internal membranes that compartmentalize their functions**

- The basic structural and functional unit of every organism is one of two types of cells: prokaryotic or eukaryotic
- Only organisms of the domains Bacteria and Archaea consist of prokaryotic cells
- Protists, fungi, animals, and plants all consist of eukaryotic cells

© 2018 Pearson Education Ltd.



## Comparing Prokaryotic and Eukaryotic Cells

- Basic features of all cells:
  - Plasma membrane
  - Semifluid substance called **cytosol**
  - Chromosomes (carry genes)
  - Ribosomes (make proteins)

© 2018 Pearson Education Ltd.

- **Prokaryotic cells** are characterized by having
  - No nucleus
  - DNA in an unbound region called the **nucleoid**
  - No membrane-bound organelles
  - **Cytoplasm** bound by the plasma membrane

© 2018 Pearson Education Ltd.







- **Eukaryotic cells** are characterized by having
  - DNA in a nucleus that is bounded by a double membrane
  - Membrane-bound organelles
  - **Cytoplasm** in the region between the plasma membrane and nucleus
- Eukaryotic cells are generally much larger than prokaryotic cells

© 2018 Pearson Education Ltd.

- The **plasma membrane** is a selective barrier that allows sufficient passage of oxygen, nutrients, and waste to service the volume of every cell


© 2018 Pearson Education Ltd.



- Metabolic requirements set upper limits on the size of cells
- The surface area to volume ratio of a cell is critical
- As a cell increases in size, its volume grows proportionately more than its surface area

© 2018 Pearson Education Ltd.

Figure 7.7




© 2018 Pearson Education Ltd.

## A Panoramic View of the Eukaryotic Cell


- A eukaryotic cell has internal membranes that divide the cell into compartments—the organelles
- The basic fabric of biological membranes is a double layer of phospholipids and other lipids
- Plant and animal cells have most of the same organelles

© 2018 Pearson Education Ltd.

Figure 7.8a




## BioFlix: Tour of an Animal Cell




© 2018 Pearson Education Ltd.

Figure 7.8b




© 2018 Pearson Education Ltd.

## BioFlix: Tour of a Plant Cell



© 2018 Pearson Education Ltd.

## Video: *Chlamydomonas*



© 2018 Pearson Education Ltd.

### **Concept 7.3: The eukaryotic cell's genetic instructions are housed in the nucleus and carried out by the ribosomes**

- The nucleus contains most of the DNA in a eukaryotic cell
- Ribosomes use the information from the DNA to make proteins

© 2018 Pearson Education Ltd.

### **The Nucleus: Information Central**

- The **nucleus** contains most of the cell's genes and is usually the most conspicuous organelle
- The **nuclear envelope** encloses the nucleus, separating it from the cytoplasm
- The nuclear envelope is a double membrane; each membrane consists of a lipid bilayer

© 2018 Pearson Education Ltd.









Figure 7.9b




© 2018 Pearson Education Ltd.

Figure 7.9c



© 2018 Pearson Education Ltd.

Figure 7.9d



Nuclear lamina (TEM)

© 2018 Pearson Education Ltd.

- Pores, lined with a structure called a pore complex, regulate the entry and exit of molecules from the nucleus
- The nuclear size of the envelope is lined by the **nuclear lamina**, which is composed of proteins and maintains the shape of the nucleus

© 2018 Pearson Education Ltd.

- In the nucleus, DNA is organized into discrete units called **chromosomes**
- Each chromosome contains one DNA molecule associated with proteins, called **chromatin**
- Chromatin condenses to form discrete chromosomes as a cell prepares to divide
- The **nucleolus** is located within the nucleus and is the site of ribosomal RNA (rRNA) synthesis

© 2018 Pearson Education Ltd.

### **Ribosomes: Protein Factories**

- **Ribosomes** are complexes made of ribosomal RNA and protein
- Ribosomes carry out protein synthesis in two locations:
  - In the cytosol (free ribosomes)
  - On the outside of the endoplasmic reticulum or the nuclear envelope (bound ribosomes)

© 2018 Pearson Education Ltd.






Figure 7.10b



© 2018 Pearson Education Ltd.

### Concept 7.4: The endomembrane system regulates protein traffic and performs metabolic functions in the cell

- The **endomembrane system** consists of
  - Nuclear envelope
  - Endoplasmic reticulum
  - Golgi apparatus
  - Lysosomes
  - Vacuoles
  - Plasma membrane
- These components are either continuous or connected via transfer by **vesicles**


© 2018 Pearson Education Ltd.

## The Endoplasmic Reticulum: Biosynthetic Factory

- The **endoplasmic reticulum (ER)** accounts for more than half of the total membrane in many eukaryotic cells
- The ER membrane is continuous with the nuclear envelope
- There are two distinct regions of ER:
  - **Smooth ER**, which lacks ribosomes
  - **Rough ER**, whose surface is studded with ribosomes

© 2018 Pearson Education Ltd.

Figure 7.11



© 2018 Pearson Education Ltd.

### ***Functions of Smooth ER***

- The smooth ER
  - Synthesizes lipids
  - Metabolizes carbohydrates
  - Detoxifies drugs and poisons
  - Stores calcium ions

© 2018 Pearson Education Ltd.

### ***Functions of Rough ER***

- The rough ER
  - Has bound ribosomes, which secrete **glycoproteins** (proteins covalently bonded to carbohydrates)
  - Distributes **transport vesicles**, secretory proteins surrounded by membranes
  - Is a membrane factory for the cell


© 2018 Pearson Education Ltd.

## The Golgi Apparatus: Shipping and Receiving Center

- The **Golgi apparatus** consists of flattened membranous sacs called cisternae
- The Golgi apparatus
  - Modifies products of the ER
  - Manufactures certain macromolecules
  - Sorts and packages materials into transport vesicles

© 2018 Pearson Education Ltd.

Figure 7.12



© 2018 Pearson Education Ltd.



## Lysosomes: Digestive Compartments

- A **lysosome** is a membranous sac of hydrolytic enzymes that can digest macromolecules
- Lysosomal enzymes work best in the acidic environment inside the lysosome
- Hydrolytic enzymes and lysosomal membranes are made by rough ER and then transferred to the Golgi apparatus for further processing

© 2018 Pearson Education Ltd.

- Some types of cell can engulf another cell by **phagocytosis**; this forms a food vacuole
- A lysosome fuses with the food vacuole and digests the molecules
- Lysosomes also use enzymes to recycle the cell's own organelles and macromolecules, a process called autophagy

© 2018 Pearson Education Ltd.

Figure 7.13







Figure 7.13a






## Animation: Lysosome Formation



## Video: Phagocytosis in Action



© 2018 Pearson Education Ltd.

## Vacuoles: Diverse Maintenance Compartments


- **Vacuoles** are large vesicles derived from the ER and Golgi apparatus
- Vacuoles perform a variety of functions in different kinds of cells

© 2018 Pearson Education Ltd.

- **Food vacuoles** are formed by phagocytosis
- **Contractile vacuoles**, found in many freshwater protists, pump excess water out of cells
- **Central vacuoles**, found in many mature plant cells, hold organic compounds and water


© 2018 Pearson Education Ltd.

Figure 7.14



© 2018 Pearson Education Ltd.

### **Video: *Paramecium* Vacuole**




© 2018 Pearson Education Ltd.

### **The Endomembrane System: *A Review***

- The endomembrane system is a complex and dynamic player in the cell's compartmental organization

© 2018 Pearson Education Ltd.

Figure 7.15



© 2018 Pearson Education Ltd.

### Concept 7.5: Mitochondria and chloroplasts change energy from one form to another

- **Mitochondria** are the sites of cellular respiration, a metabolic process that uses oxygen to generate ATP
- **Chloroplasts**, found in plants and algae, are the sites of photosynthesis
- Peroxisomes are oxidative organelles

© 2018 Pearson Education Ltd.

## The Evolutionary Origins of Mitochondria and Chloroplasts


- Mitochondria and chloroplasts have similarities with bacteria:
  - Enveloped by a double membrane
  - Contain free ribosomes and circular DNA molecules
  - Grow and reproduce somewhat independently in cells
- These similarities led to the **endosymbiont theory**

© 2018 Pearson Education Ltd.

- The endosymbiont theory suggests that an early ancestor of eukaryotes engulfed an oxygen-using nonphotosynthetic prokaryotic cell
- The engulfed cell formed a relationship with the host cell, becoming an endosymbiont
- The endosymbionts evolved into mitochondria
- At least one of these cells may have then taken up a photosynthetic prokaryote, which evolved into a chloroplast

© 2018 Pearson Education Ltd.









## Mitochondria: Chemical Energy Conversion

- Mitochondria are found in nearly all eukaryotic cells
- They have a smooth outer membrane and an inner membrane folded into **cristae**
- The inner membrane creates two compartments: intermembrane space and **mitochondrial matrix**
- Some metabolic steps of cellular respiration are catalyzed in the mitochondrial matrix
- Cristae present a large surface area for enzymes that synthesize ATP

© 2018 Pearson Education Ltd.






## Chloroplasts: Capture of Light Energy

- Chloroplasts contain the green pigment chlorophyll, as well as enzymes and other molecules that function in photosynthesis
- Chloroplasts are found in leaves and other green organs of plants and in algae

© 2018 Pearson Education Ltd.

Figure 7.18



© 2018 Pearson Education Ltd.

